

Endpoint dilution using an antibody to the adenoviral hexon protein may be used to quantitate virus production or infection efficiency of target cells (Armentano et al. (1995) Hum. Gene Ther. 6:1343-53.

In addition to containing the DNA sequences encoding one or more transgenes, the chimeric adenoviral vectors of the invention may contain any expression control sequences such as a promoter or enhancer, a polyadenylation element, and any other regulatory elements that may be used to modulate or increase expression, all of which are operably linked in order to allow expression of the transgene. The use of any expression control sequences, or regulatory elements, which facilitate expression of the transgene is within the scope of the invention. Such sequences or elements may be capable of generating tissue-specific expression or be susceptible to induction by exogenous agents or stimuli.

#### **Construction of Retroviral Vectors**

The antigen presenting cells described herein can also be genetically modified with retroviral vectors produced recombinantly by procedures already taught in the art. For example, WO 94/29438 describes the construction of retroviral packaging plasmids and packaging cell lines. As is apparent to the skilled artisan, the retroviral vectors useful in the methods of this invention are capable of infecting the cells described herein. The techniques used to construct vectors, and transfect and infect cells are widely practiced in the art. Examples of retroviral vectors are those derived from murine, avian or primate retroviruses. Retroviral vectors based on the Moloney murine leukemia virus (MoMLV) are the most commonly used because of the availability of retroviral variants that efficiently infect human cells. Other suitable vectors include those based on the Gibbon Ape Leukemia Virus (GALV) or HIV.

The range of host cells that may be infected by a retrovirus or retroviral vector is determined by the viral envelope protein. The recombinant virus can be used to infect virtually any other cell type recognized by the env protein provided by the packaging cell, resulting in the integration of the viral genome in the

transduced cell and the stable production of the foreign gene product. In general, murine ecotropic env of MoMLV allows infection of rodent cells, whereas amphotropic env allows infection of rodent, avian and some primate cells, including human cells. Amphotropic packaging cell lines for use with MoMLV systems are known in the art and commercially available and include, but are not limited to, PA12 and PA317. Miller et al. (1985) Mol. Cell. Biol. 5:431-437; Miller et al. (1986) Mol. Cell. Biol. 6:2895-2902; and Danos et al. (1988) Proc. Natl. Acad. Sci. USA 85:6460-6464. Xenotropic vector systems exist which also allow infection of human cells.

The host range of retroviral vectors has been altered by substituting the env protein of the base virus with that of a second virus. The resulting, "pseudotyped", virus has the host range of the virus donating the envelope protein and expressed by the packaging cell line. Recently, the G-glycoprotein from vesicular stomatitis virus (VSV-G) has been substituted for the MoMLV env protein. Burns et al. (1993) Proc. Natl. Acad. Sci USA 90:8033-8037; and PCT patent application WO 92/14829. Since infection is not dependent on a specific receptor, VSV-G pseudotyped vectors have a broad host range.

Usually, the vectors will contain at least two heterologous genes or gene sequences: (i) the therapeutic gene to be transferred; and (ii) a marker gene that enables tracking of infected cells.

The viral vector can be modified to incorporate chimeric envelope proteins or nonviral membrane proteins into retroviral particles to improve particle stability and expand the host range or to permit cell type-specific targeting during infection. The production of retroviral vectors that have altered host range is taught, for example, in WO 92/14829 and WO 93/14188. Retroviral vectors that can target specific cell types *in vivo* are also taught, for example, in Kasahara et al. (1994) Science 266:1373-1376. Kasahara et al. describe the construction of a Moloney leukemia virus (MoMLV) having a chimeric envelope protein consisting of human erythropoietin (EPO) fused with the viral envelope protein. This hybrid virus shows tissue tropism for human red blood progenitor cells that bear the

receptor for EPO, and is therefore useful in gene therapy of sickle cell anemia and thalassemia. Retroviral vectors capable of specifically targeting infection of cells are preferred for *in vivo* gene therapy.

5       The vector construct is introduced into a packaging cell line which will generate infectious virions. Packaging cell lines capable of generating high titers of replication-defective recombinant viruses are known in the art, see for example, WO 94/29438. Viral particles are harvested from the cell supernatant and purified for *in vivo* infection using methods known in the art such as by filtration of supernatants 48 hours post transfection. The viral titer is determined by infection  
10       of a constant number of appropriate cells (depending on the retrovirus) with titrations of viral supernatants. The transduction efficiency can be assayed 48 hours later by a variety of methods, including Southern blotting.

      After viral transduction, the presence of the viral vector in the transduced cells or their progeny can be verified such as by PCR. PCR can be performed to  
15       detect the marker gene or other virally transduced sequences. Generally, periodic blood samples are taken and PCR conveniently performed using e.g. NeoR probes if the NeoR gene is used as marker. The presence of virally transduced sequences in bone marrow cells or mature hematopoietic cells is evidence of successful reconstitution by the transduced cells. PCR techniques and reagents are well  
20       known in the art, See, generally, PCR PROTOCOLS, A GUIDE TO METHODS AND APPLICATIONS. Innis, Gelfand, Sninsky & White, eds. (Academic Press, Inc., San Diego, 1990) and commercially available (Perkin-Elmer).

*In vitro/ex vivo*, exposure of human DCs to adenovirus (Ad) vector at a multiplicity of infection (MOI) of 500 for 16-24 h in a minimal volume of serum-free medium reliably gives rise to transgene expression in 90-100% of DCs. The  
25       efficiency of transduction of DCs or other APCs can be assessed by immunofluorescence using fluorescent antibodies specific for the tumor antigen being expressed (Kim et al., 1997). Alternatively, the antibodies can be conjugated to an enzyme (e.g. HRP) giving rise to a colored product upon reaction

with the substrate. The actual amount of antigen being expressed by the APCs can be evaluated by ELISA.

Transduced APCs can subsequently be administered to the host via an intravenous, subcutaneous, intranasal, intramuscular or intraperitoneal route of delivery.

*In vivo* transduction of DCs, or other APCs, can potentially be accomplished by administration of Ad (or other viral vectors) via different routes including intravenous, intramuscular, intranasal, intraperitoneal or cutaneous delivery. The preferred method is cutaneous delivery of Ad vector at multiple sites using a total dose of approximately  $1 \times 10^{10}$ - $1 \times 10^{12}$  i.u. Levels of *in vivo* transduction can be roughly assessed by co-staining with antibodies directed against APC marker(s) and the TAA being expressed. The staining procedure can be carried out on biopsy samples from the site of administration or on cells from draining lymph nodes or other organs where APCs (in particular DCs) may have migrated. The amount of TAA being expressed at the site of injection or in other organs where transduced APCs may have migrated can be evaluated by ELISA on tissue homogenates.

Although viral gene delivery is more efficient, DCs can also be transduced *in vitro/ex vivo* by non-viral gene delivery methods such as electroporation, calcium phosphate precipitation or cationic lipid/plasmid DNA complexes. Transduced APCs can subsequently be administered to the host via an intravenous, subcutaneous, intranasal, intramuscular or intraperitoneal route of delivery.

*In vivo* transduction of DCs, or other APCs, can potentially be accomplished by administration of cationic lipid/plasmid DNA complexes delivered via the intravenous, intramuscular, intranasal, intraperitoneal or cutaneous route of administration. Gene gun delivery or injection of naked plasmid DNA into the skin also leads to transduction of DCs. Intramuscular delivery of plasmid DNA may also be used for immunization.

The transduction efficiency and levels of transgene expression can be assessed as described above for viral vectors.

### Expansion of Immune Effector Cells

5           The present invention makes use of these APCs to stimulate production of an enriched population of antigen-specific immune effector cells. The antigen-specific immune effector cells are expanded at the expense of the APCs, which die in the culture. The process by which naïve immune effector cells become educated by other cells is described essentially in Coulie (1997) *Molec. Med.*  
10       Today 3:261-268.

          In a preferred embodiment, the antigen-specific immune effector cells are CTLs. In one aspect, the cytotoxic T cells are polyclonal T cells isolated from a site of cytotoxic T cell infiltration from a subject. Alternatively, such cells may be isolated from a site of cytotoxic T cell infiltration from two or more subjects or  
15       human patients, in which the subjects share an MHC halotype. In another embodiment, the CTLs may be two or more cytotoxic T cell lines. In yet another embodiment, the CTLs may be any combination of the foregoing.

          In a further aspect of the invention, the site of cytotoxic T cell infiltration is a tumor. The tumors from which cells or cell lines are obtained can be the same  
20       type of tumor in different individuals with a shared MHC halotype or different types of tumors from different subjects who share an MHC haplotype.

          The APCs prepared as described above are mixed with naïve immune effector cells. Preferably, the cells may be cultured in the presence of a cytokine, for example IL2. Because dendritic cells secrete potent immunostimulatory  
25       cytokines, such as IL2, it may not be necessary to add supplemental cytokines during the first and successive rounds of expansion. In any event, the culture conditions are such that the antigen-specific immune effector cells expand (*i.e.* proliferate) at a much higher rate than the APCs. Multiple infusions of APCs and optional cytokines can be performed to further expand the population of antigen-  
30       specific cells.

In one embodiment, the immune effector cells are T cells and are specific for tumor-specific antigens which are presented by the APCs.

### **Compositions**

5           This invention also provides compositions containing any of the above-mentioned proteins, muteins, polypeptides, nucleic acid molecules, vectors, cells antibodies and fragments thereof, and an acceptable solid or liquid carrier. When the compositions are used pharmaceutically, they are combined with a "pharmaceutically acceptable carrier" for diagnostic and therapeutic use. These  
10       compositions also can be used for the preparation of medicaments for the diagnosis and treatment of diseases such as cancer.

### **Tumor Protection in Animal Models**

Applicants are the first to establish that, based on the animal models  
15       described below, prevaccination with the compositions of this invention will prevent or delay onset of disease.

The murine B16 melanoma model was used. In this model, C57BL/6 mice were immunized with bone marrow-derived DCs transduced with an Ad vector encoding either human gp100 (Ad/hugp100) or mouse gp100 (Ad/mgpl00). Mice  
20       immunized against heterologous human gp100 developed a protective immune response and were resistant to a lethal subcutaneous challenge of B16 melanoma cells (syngeneic tumor cell line that expresses gp100). In contrast, mice immunized with homologous mouse gp100 failed to mount a protective immune response against B16 melanoma cells and developed tumors at the site of B16 cell  
25       injection. This finding illustrates the difficulty in breaking tolerance against a self antigen (mouse gp100). The corresponding heterologous antigen from a different species (human gp100), however, is likely to contain several Class I and Class II-associated epitopes that will be recognized as foreign and elicit CD8<sup>+</sup> and CD4<sup>+</sup> T cell responses, respectively. The induction of cross-reactive CTLs that recognize

both the heterologous and homologous self-antigen can then lead to lysis of host tumor cells.

Unfortunately, this type of animal model cannot be used to test the efficacy of modified or heterologous tumor antigens being considered for use in humans since mice and humans recognize different epitopes, primarily as a result of differences in their MHC molecules. It may be possible, however, to use the allogeneic human peripheral blood lymphocyte - severe combined immunodeficiency mouse (Hu-PBL-SCID) model. SCID mice lack mature B and T lymphocytes and can be reconstituted with human PBLs. It may be possible to immunize such mice with test antigen to induce a response in adoptively transferred human PBLs and evaluate protection against challenge with a human tumor cell line (Mosier et al., 1988; Parney et al., 1997; Albert et al., 1997).

Another possibility is immunization of HLA-A2.1 transgenic mice to reproduce the immune reactivity of HLA-A2 individuals.

#### **Adoptive Immunotherapy and Vaccines**

The expanded populations of antigen-specific immune effector cells of the present invention find use in adoptive immunotherapy regimes and as vaccines.

Adoptive immunotherapy methods involve, in one aspect, administering to a subject a substantially pure population of educated, antigen-specific immune effector cells made by culturing naïve immune effector cells with APCs as described above. Preferably, the APCs are dendritic cells.

In one embodiment, the adoptive immunotherapy methods described herein are autologous. In this case, the APCs are made using parental cells isolated from a single subject. The expanded population also employs T cells isolated from that subject. Finally, the expanded population of antigen-specific cells is administered to the same patient.

In another embodiment, the adoptive immunotherapy methods are allogeneic. Here, cells from two or more patients are used to generate the APCs, and stimulate production of the immune effector cells. For instance, cells from

other healthy or diseased subjects can be used to generate antigen-specific cells in instances where it is not possible to obtain autologous T cells and/or dendritic cells from the subject providing the biopsy. The expanded population can be administered to any one of the subjects from whom cells were isolated, or to  
5 another subject entirely.

In a further embodiment, APCs or immune effector cells are administered with an effective amount of a stimulatory cytokine, such as IL-2 or a co-stimulatory molecule.

The agents identified herein as effective for their intended purpose can be administered to subjects or individuals susceptible to or at risk of developing a  
10 disease, such as cancer. When the agent is administered to a subject such as a mouse, a rat or a human patient, the agent can be added to a pharmaceutically acceptable carrier and systemically or topically administered to the subject. To determine patients that can be beneficially treated, a tumor regression can be  
15 assayed. Therapeutic amounts can be empirically determined and will vary with the pathology being treated, the subject being treated and the efficacy and toxicity of the therapy. When delivered to an animal, the method is useful to further confirm efficacy of the agent. As an example of an animal model, groups of nude mice (Balb/c NCR nu/nu female, Simonsen, Gilroy, CA) are each subcutaneously  
20 inoculated with about  $10^5$  to about  $10^9$  hyperproliferative, cancer or target cells as defined herein. When the tumor is established, the agent is administered, for example, by subcutaneous injection around the tumor. Tumor measurements to determine reduction of tumor size are made in two dimensions using vernier calipers twice a week. Other animal models may also be employed as  
25 appropriate.

Administration *in vivo* can be effected in one dose, continuously or intermittently throughout the course of treatment. Methods of determining the most effective means and dosage of administration are well known to those of skill in the art and will vary with the composition used for therapy, the purpose of  
30 the therapy, the target cell being treated, and the subject being treated. Single or



multiple administrations can be carried out with the dose level and pattern being selected by the treating physician. Suitable dosage formulations and methods of administering the agents can be found below.

5 The agents and compositions of the present invention can be used in the manufacture of medicaments and for the treatment of humans and other animals by administration in accordance with conventional procedures, such as an active ingredient in pharmaceutical compositions.

10 More particularly, an agent of the present invention also referred to herein as the active ingredient, may be administered for therapy by any suitable route including nasal, topical (including transdermal, aerosol, buccal and sublingual), parenteral (including subcutaneous, intramuscular, intravenous and intradermal) and pulmonary. It will also be appreciated that the preferred route will vary with the condition and age of the recipient, and the disease being treated.

15 The compositions and methods described herein are particularly useful in providing or inducing a prophylactic immune response in an animal. Animals in a pre-disease state or in a disease free interval, i.e., having or pre-disposed to a condition subject to immune surveillance, are most suitably treated by the methods and compositions described herein. Such conditions involve the activation of an immune response in a diseased state or period.

20

#### Example

25 Dendritic cells were derived from peripheral blood of a HLA-A2+ human donor using standard GM-CSF/IL-4 culture technique. After six days, the cells were infected with an adenovirus (serotype 2) construct encoding human gp 100. At T=7 days, they were restimulated with  $5 \times 10^5$  infected autologous DCs (from frozen stocks) and given 50 U/ml rhIL-2. At T=14 days, the cultures were treated with leucyl-leucyl-methyl ester in order to eliminate NK cell activity. Immediately after treatment, the cells were washed thoroughly and replated along  
30 with  $1 \times 10^7$  mitomycin C-treated autologous PBMC as feeders. At T=21 days,

cells were split and replated at  $5 \times 10^5$  cells/ml in Iscoves/10% human AB serum/1000U rhIL-2/ml. At T=26 days, the CTLs raised against gp100 were tested in  $^{51}\text{Cr}$ -release assay using peptide pulsed T2 cells as targets. Effector CTLs were TIL 1520 which specifically recognize an HLA-A-A2 restricted epitope of the wild-type human gp100 protein. Figure 5 shows the results of this assay. There is epitope-specific recognition that was not present prior to education of the T cells with the infected DCs.

It is to be understood that while the invention has been described in conjunction with the above embodiments, that the foregoing description and the following examples are intended to illustrate and not limit the scope of the invention. For example, any of the above-noted compositions and/or methods can be combined with known therapies or compositions. Other aspects, advantages and modifications within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

CLAIMS

1. A method of inducing a prophylactic immune response to a self-antigen in a subject, comprising administering to the subject an effective amount  
5 of the antigen or an altered form of the antigen.

2. The method of claim 1, wherein the antigen is administered as a polynucleotide coding for the self-antigen.

10 3. The method of claim 2, wherein the polynucleotide is delivered as naked DNA.

4. The method of claim 2, wherein the polynucleotide is delivered in a gene delivery vehicle.

15 5. The method of claim 1, further comprising administering an effective amount of an immunostimulatory agent to the subject.

20 6. The method of claim 5, wherein the immunostimulatory agent is administered as a polynucleotide coding for the immunostimulatory agent.

7. The method of claim 1, wherein the antigen is administered in an antigen presenting cell.

25 8. The method of claim 7, wherein the antigen presenting cell has been genetically modified by insertion of a polynucleotide coding for the antigen.

30 9. The method of claim 7, wherein the antigen presenting cell is a foster antigen presenting cell, a hybrid antigen presenting cell, or a pulsed antigen presenting cell.

10. The method of claim 7, wherein the antigen presenting cell is a dendritic cell.

5 11. The method of claim 7, further comprising administering an effective amount of an immunostimulatory agent to the subject.

12. The method of claim 11, wherein the immunostimulatory agent is administered as a polynucleotide coding for the immunostimulatory agent.

10 13. The method of claim 1 or 7, wherein the self-antigen is a tumor associated antigen (TAA).

14. A method of providing a prophylactic immune response to a self-antigen in a subject, comprising administering to the subject an effective amount of educated immune effector cells, educated to specifically recognize and lyse cells expressing the self-antigen or an altered form of the self-antigen.

15 16. The method of claim 14, wherein the immune effector cells have been produced by stimulating naïve immune effector cells with antigen presenting cells that present the antigen or an altered self-antigen to the naïve immune effector cells.

20 17. The method of claim 14, wherein the educated immune effector cells are produced *ex vivo*.

25 18. The method of claim 14, wherein the educated immune effector cells are produced *in vivo*.

18. The method of claim 14, further comprising administering an effective amount of an immunostimulatory agent.

5 19. The method of claim 18, wherein the immunostimulatory agent is administered as a polynucleotide coding for the immunostimulatory agent.

20. The method of claim 1 or 14, wherein the subject is characterized as being in a disease-free state but genetically predisposed to a condition subject to immune surveillance.

10


21. The method of claim 20, wherein the condition is associated with the presence of the HER-2/neu gene in the subject.

22. The method of claim 1 or 14, wherein the subject is characterized as being in a disease free interval of a condition subject to immune surveillance.

15

23. The method of claim 22, wherein the condition is melanoma.

1/9



HUMAN	1	MDLVLKRCLL	HLAVIGALLA	VGATKVPRNQ	DWLGVSQRLR	TKAWNRLQLYP
MOUSE		--*G-QR-SF-	PLV-LSA---	---LEGS---	-----P---V	--T-----
	51	EWTEAQRDC	WRGCGVSLKV	SNDGPTLIGA	NASFSLALNF	PGSQKVLPDG
		----V-GSN-	-----R-	I----- --	-----H-	-----
	101	QVIWVNNTII	NGSQVWGGQP	VYPQETDDAC	IFPDGGPCPS	GSWSQKRFSV
		----A----	-----	---P----	V-----	-PKPP----
	151	YVWKIWGQYW	QVLGCPVSGL	SIGTGRAMLG	THIMEVTVYH	RRGSRSYVPL
		-----K--	-----R-	--A--H-K--	-----	----Q----
	201	AHSSSAFTIT	DQVPFVSVS	QLRALDGGNK	HFLRNQPLTF	ALQLHDPSCY
		--A--T----	-----	--Q-----ET-	-----H--I-	-----
	251	LAEADLSYTW	DFGDSSGTLI	SRALVVTHTY	LEPGPVTAQV	VLQAAIPLTS
		-----GT----	-----D----	-----S-S-	-----V-	-----

FIG. 1A

2/9

```

301 CGSSPVPGIT DGHRTAEAP NITAGQVPTT EVVGTTPGQA PTAEPSGITS
    ----- --YM----- G--SR-GT-- -----M--TQ-----V

351 VQVPTTEVIS TAPVQMPTAE STGMTPEKVP VSEVMGTILA EMSTPEATGM
    --M-----TA -TSE--L--* ***** **A-ID----- -V--T-G--T

401 TPAEVSIVVL SGTIAAQVTT TEWVETTARE LPIPEPEGPD ASSIMSTESI
    --T*****P -----V--A-- --***** **----- --PLLP-Q-S

451 TGGLGPLLDG TATRLVKRQ VPLDCVLYRY GSFSVTLDIV QGIESAEILQ
    ---IS-----D -D-IM----- -----LA-----

501 AVPSGEGDAF ELTVSCQGGL PKEACMEISS PGCQPPAQR L CQVLPSPAC
    ---FS----- -----D----- -----S-P---D-

551 QLVLHQILKG GSGTYCLNVS LADTNSLAVV STQLIMPQE AGLGQVPLIV
    -----V----- -----A-----A-----VV---D G---A--L-

601 GILLVLMVV LASLIYRRRL MKQDFVLPQL PHSSHVLR L PRIFCSCPIG
    -----V----- -----H-H-- K--G*--S-M --G-T----- -PV-RARGL-

651 ENSPLLSGQQ VX
    -----

```

FIG. 1B

3/9




FIG. 2A

A	mMART	10 20 30 40 50	ATGCCCCAAGAAGACATTCACTT-----TGGTTATCCCAGGAAGGGCACAGACGCTCC	30 40 50
	hMART	10 20 30 40 50 60	ATGCCCCAAGAAGATGCTCACTTTCATCTATGTTACCCCAAGAAGGGCACGGCCACTCT	30 40 50 60
	mMART	60 70 80 90 100 110	TATGTCACCTGCTGAAGAGCGCCGACGGATCGGCATCCTGATCGTGGTCCCTGGGATTGCT	90 100 110
	hMART	70 80 90 100 110 120	TACACCACGGCTGAAGAGCGCCGCTGGGATCGGCATCCTGACAGTGTATCCCTGGGAGTCTTA	90 100 110 120
	mMART	120 130 140 150 160 170	CTGCTTATCGGCTGCTGGTACTGTAGAAGACGAAGTGGATACAGAACCTTGTATGGACAAA	130 140 150 160 170
	hMART	130 140 150 160 170 180	CTGCTCATCGGCTGTGGTATTGTAGAAGACGAATGGATACAGAGCCTTGTATGGATAAA	130 140 150 160 170 180
	mMART	180 190 200 210 220 230	AGCGCTCATATTGGTATTCAAAAACCTCAAGGGAAGATGCTCATGTGAGAGCCCTGAT	190 200 210 220 230
	hMART	190 200 210 220 230 240	AGCTTTCATGTTGGCACTCAATGTGCCCTTAACAAGAAGATGCCCAACAAGAAGGGTTTGT	190 200 210 220 230 240



**B**

mMART	240	250	260	270	280	290
	CACCAGGACAGCCGAC	TGCTCTTCTCAAGAGAAATCCCA	TGAGCCCGTGGTTC	CCCAACGCT		
hMART	250	260	270	280	290	300
	CATCGGACAGCAAGTG	CTCTTCAAGAGAAAACTG	GAACCTGIGGTTCCCA	ATGCT		
mMART	300	310	320	330	340	
	CCGCTGCCCTATGAGA	AGCTCTCT-----TCACCGCCACCTT	ATTCACCCCTGA			
hMART	310	320	330	340	350	
	CCACCTGCTTATGAGAA	ACTCTCTGCAGAACAGTCA	CCACCACCTTATTCACCTTAA			
mMART	10	20	30	40	50	
	MPQEDIHF--GYPRKG	RRRSYVTAEAAAGIGILIV	LGIALIGCWYCRRRSGYRT	LMDK		
hMART	10	20	30	40	50	60
	MPREDAHF	IYGYPKKGHGSYTTAEAA	GIGILTVILGVLLIGCWYCRRR	NGYRALMDK		
mMART	60	70	80	90	100	110
	RRHIGIQKTSRERC	SCSPDHQDSRLSSQEKSHQ	VPVNAPPAYEKL	SPPPYSPX		
hMART	70	80	90	100	110	
	SLHVG	TQCALIRRC	PQEGFDRDSKVS	LQKNC	EPVVPNAPPAYEKL	SAEQSPPPYSPX

FIG. 2B

5/9

FIG. 3

FIG. 3A

FIG. 3B

```

1  AGCAGACAGAGGACTCTCATTAAGGAAGG  TGTCCTGTGCCCTTGACCCCTACAAGATGCCA
MetPro

120  ACGGCTGAAGAGGCCGCTGGGATCGGCATC  CIGACAGTGAATCCITGGGAGTCTTACTGCTC
23  ThrAlaGluGluAlaAlaGlyIleGlyIle  LeuThrValIleLeuGlyValLeuLeuLeu

240  CATGTTGGCACITCAATGTGCCITTAACAAGA  AGATGCCCCACAAGAAGGTTTGTATCATCGG
63  HisValGlyThrGlnCysAlaLeuThrArg  ArgCysProGlnGluGlyPheAspHisArg

360  GCATTATGAGAAACTCTCTGCAGAACAGTCA  CCACCACCITATTACCTTAAGAGCCACGG
103  AlaTyrGluLysLeuSerAlaGluGlnSer  ProProProTyrSerPro

480  ATCTAATGTTCTCCTTTGGAAATGGGTAGG  AAAAATGCAAGCCATCCTCTAATAATAAGTC
600  TATTAAATTGGGAAAACTCCATCAATAAAT  GTTGCAATGCATGATACTATCTGTGCCAGA
720  GGGGCCATCCAAATTTCTCTTTTACTTGAAAT  TTGGCTAATAACAACAACCTAGTCAGGTTTTCG
840  GATACITTTACAGGTTAAGACAAAGGGTTG  ACTGGCCTATTATCTGATCAAGAACAATGT
960  CTATAGCTCTTTTTTTTGGAGATGGAGTTT  CGCTTTTGTGGCCAGGCTGGAGTGCAATG
1080  CCTCCTGAGTAGCTGGGATTACAGGCGGCG  GCCACTATGCCCTGACTAATTTGTAGTTT
1200  TCTGCCCCGCTCAGCCTCCCAAAGTCTGG  AATTACAGGCGTGAGCCACCACGCTGGCT
1320  AATGCTATTCTAACTAATGACAAAGTATTT  CTACTAAACCAGAAATTGGTAGAAGGATTT
1440  TACCTATGGCAATTAGCTCTCTTGGGTTC  CCAAATCCCTCTCACAAAGATGTGCAGAAG

```

FIG. 3A

6/9

AGAGAAAGATGCTCACCATTATGTTAC	CCCAAGAAGGGGCACGGCCACTCTACACC	119
ArgGluAspAlaHisPheIleTyrGlyTyr	ProLysLysGlyHisGlyHisSerTyrThr	22
ATCGGCTGTGGTATTGTAGAAGACGAAAT	GGATACAGAGCCCTTGATGGATAAAAGTCTT	239
<u>IleGlyCysTrpTyrCysArgArgArgAsn</u>	GlyTyrArgAlaLeuMetAspLysSerLeu	62
GACAGCAAAGTGTCCTCTTCAAGAGAAAAAC	TGTGAACCTGTGGTTCCCAATGCTCCACCT	359
AspSerLysValSerLeuGlnGluLysAsn	CysGluProValValProAsnAlaProPro	102
AGACACCTGAGACATGCTGAAATTATTCT	CTCACACCTTTTGGCTTGAATTTAATACAGAC	479
		118
AGTGTTAAAAATTTIAGTAGGTCGGCTAGCA	GTACTAATCATGTGAGGAAATGATGAGAAA	599
GGTAAATGTTAGTAAATCCATGGTGTTATTT	TCTGAGAGACAGCAATTCAAGTGGGTATTCT	719
AACCTTGACCGACATGAACGTGTACACAGAA	TTGTTCCAGTACTATGGAGTGTCACAAAG	839
CAGCAATGTCCTTTGTGCTCTAAAATTCT	ATTATACTACAATAATATATTGTAAAGATC	959
GC GGATCTTTGGCTCACCATAACTCCGCC	TCCCAGGTTCAAGCAATTCCTCCCTTAG	1079
AGTAGAGACCGGGTTCTCCATGTGGTCA	GGCTGGTCTCAAACTCCTGACCICAGGTGA	1199
GGATCCCTATATCTTAGGTAAGACATAAATAC	GCAGTCTAATTACATTTACACTTCAAGGCTC	1319
AAATAAGTAAAGCTACTATGTACTGCCIT	AGTGTGATGCTGTGCTACTGCCITAAATG	1439
AAATCATAAAGGATCAGAGATTCTGAAAAA	AAAAAAAATAAAAAATAAAAAATAAAAAA	1559

FIG. 3B

FIG. 4A
FIG. 4B

FIG. 4

MOUSE TRP2

```

1  MGLVGWGLLL GCLGCGILLR ARAQFPRVCM TLDGVLNKEC CPPLGPEATN
51  ICGFLEGRGQ CAEVQTDTRP WSGPYILRNQ DDREQWPRKF FNRCKCTGN
101 FAGYNCGGCK FGWTGPDCNR KKPAILRRNI HSLTAQEREQ FLGALDLAKK
151 SIHPDYVITT QHWLGLLGN GTQPQIANCE VYDFFVWLHY YSVRDILLGP
201 GRPYKAIDFS HQGPAFVTWH RYHLLWLERE LQRLTGNESF ALPYWNFATG
251 KNECDVCTDD WLGAARQDDP TLISRNSRFS TWEIVCDSD DYNRRVTLCN
301 GTYEGLLRRN KVGRNNEKLP TLKNVQDCLS LQKFDSPFF QNSTFSFRNA
351 LEGFDKADGT LDSQVMNLHN LAHSFLNGTN ALPHSAANDP VFVVLHSFTD
401 AIFDEWLKRN NPSTDWPQE LAPIGHNRMV NMVPFFPPVT NEELFLTAEQ
451 LGYNYAVDLS EEEAPVWSTT LGVVIGILGA FVLLGLLAF LQYRRLRKGY
501 APLMETGLSS KRYTEEA
    
```

FIG. 4A

80/9

## HUMAN TRP2

1 MSPLWWGFLI SCLGCKILPG AQGFPRVCM TVDSLVNKEC CPRLGAESAN  
51 VCGSQQGRGQ CTEVRADTRP WSGPYILRNQ DDRELWPRKF FHRTCKCIGN  
101 FAGYNCGDCK FGWTGPNCEK KKPPVIRQNI HSLSPQEREQ FLGALDLAKK  
151 RVHPDYVITT QHWLGLLGN GTQPQFANCS VYDFFVWLHY YSVRDTLLGP  
201 GRPYRAIDFS HQGPAFVTVH RYHLLCLERD LQRLIGNESF ALPYWNFATG  
251 RNECDVCTDQ LFGAARPDDP TLISRSRFS SWETVCDSD DYNHLVTLCN  
301 GTYEGLLRN QMGRNSMKLP TLKDIRDCLS LQKFDNPPFF QNSTFSFRNA  
351 LEGFDKADGT LDSQVMSLHN LVHSFLNGTN ALPHSAANDP IFVVLHSFTD  
401 AIFDEWMKRF NPPADAWPQE LAPIGHNRMV NMVPFFPPVT NEELFLTSDQ  
451 LGYSYAIDL PVSVEETPGWP TLLVVMGTL VALVGLFVLL AFLQYRRLRK  
501 GYTPLMETHL SSKRYTEEA

FIG. 4B

9/9

Assay of CTLs Generated From  
Normal Donor PBL With Ad-GP100-  
Infected Dendritic Cells  
(Targets=T2 Cells Pulsed with GP100-  
F9 Peptide)




FIG. 5

## SEQUENCE LISTING

<110> Nicolette, Charles A.  
Genzyme Corporation

<120> COMPOSITIONS AND METHODS FOR ANTIGEN-SPECIFIC  
VACCINATION

<130> 159792001040

<140> Unassigned

<141> 1999-03-19

<150> 60/078,890

<151> 1998-03-20

<160> 17

<170> PatentIn Ver. 2.0

<210> 1

<211> 662

<212> PRT

<213> Homo sapiens

<400> 1

Met Asp Leu Val Leu Lys Arg Cys Leu Leu His Leu Ala Val Ile Gly  
1 5 10 15

Ala Leu Leu Ala Val Gly Ala Thr Lys Val Pro Arg Asn Gln Asp Trp  
20 25 30

Leu Gly Val Ser Arg Gln Leu Arg Thr Lys Ala Trp Asn Arg Gln Leu  
35 40 45

Tyr Pro Glu Trp Thr Glu Ala Gln Arg Leu Asp Cys Trp Arg Gly Gly  
50 55 60

Gln Val Ser Leu Lys Val Ser Asn Asp Gly Pro Thr Leu Ile Gly Ala  
65 70 75 80

Asn Ala Ser Phe Ser Ile Ala Leu Asn Phe Pro Gly Ser Gln Lys Val  
85 90 95

Leu Pro Asp Gly Gln Val Ile Trp Val Asn Asn Thr Ile Ile Asn Gly  
100 105 110

Ser Gln Val Trp Gly Gly Gln Pro Val Tyr Pro Gln Glu Thr Asp Asp  
115 120 125

Ala Cys Ile Phe Pro Asp Gly Gly Pro Cys Pro Ser Gly Ser Trp Ser  
130 135 140

Gln Lys Arg Ser Phe Val Tyr Val Trp Lys Thr Trp Gly Gln Tyr Trp  
145 150 155 160

Gln Val Leu Gly Gly Pro Val Ser Gly Leu Ser Ile Gly Thr Gly Arg  
165 170 175

Ala Met Leu Gly Thr His Thr Met Glu Val Thr Val Tyr His Arg Arg  
180 185 190

Gly Ser Arg Ser Tyr Val Pro Leu Ala His Ser Ser Ser Ala Phe Thr  
 195 200 205  
 Ile Thr Asp Gln Val Pro Phe Ser Val Ser Val Ser Gln Leu Arg Ala  
 210 215 220  
 Leu Asp Gly Gly Asn Lys His Phe Leu Arg Asn Gln Pro Leu Thr Phe  
 225 230 235 240  
 Ala Leu Gln Leu His Asp Pro Ser Gly Tyr Leu Ala Glu Ala Asp Leu  
 245 250 255  
 Ser Tyr Thr Trp Asp Phe Gly Asp Ser Ser Gly Thr Leu Ile Ser Arg  
 260 265 270  
 Ala Leu Val Val Thr His Thr Tyr Leu Glu Pro Gly Pro Val Thr Ala  
 275 280 285  
 Gln Val Val Leu Gln Ala Ala Ile Pro Leu Thr Ser Cys Gly Ser Ser  
 290 295 300  
 Pro Val Pro Gly Thr Thr Asp Gly His Arg Pro Thr Ala Glu Ala Pro  
 305 310 315 320  
 Asn Thr Thr Ala Gly Gln Val Pro Thr Thr Glu Val Val Gly Thr Thr  
 325 330 335  
 Pro Gly Gln Ala Pro Thr Ala Glu Pro Ser Gly Thr Thr Ser Val Gln  
 340 345 350  
 Val Pro Thr Thr Glu Val Ile Ser Thr Ala Pro Val Gln Met Pro Thr  
 355 360 365  
 Ala Glu Ser Thr Gly Met Thr Pro Glu Lys Val Pro Val Ser Glu Val  
 370 375 380  
 Met Gly Thr Thr Leu Ala Glu Met Ser Thr Pro Glu Ala Thr Gly Met  
 385 390 395 400  
 Thr Pro Ala Glu Val Ser Ile Val Val Leu Ser Gly Thr Thr Ala Ala  
 405 410 415  
 Gln Val Thr Thr Thr Glu Trp Val Glu Thr Thr Ala Arg Glu Leu Pro  
 420 425 430  
 Ile Pro Glu Pro Glu Gly Pro Asp Ala Ser Ser Ile Met Ser Thr Glu  
 435 440 445  
 Ser Ile Thr Gly Ser Leu Gly Pro Leu Leu Asp Gly Thr Ala Thr Leu  
 450 455 460  
 Arg Leu Val Lys Arg Gln Val Pro Leu Asp Cys Val Leu Tyr Arg Tyr  
 465 470 475 480  
 Gly Ser Phe Ser Val Thr Leu Asp Ile Val Gln Gly Ile Glu Ser Ala  
 485 490 495  
 Glu Ile Leu Gln Ala Val Pro Ser Gly Glu Gly Asp Ala Phe Glu Leu  
 500 505 510



Thr Val Ser Cys Gln Gly Gly Leu Pro Lys Glu Ala Cys Met Glu Ile  
 515 520 525  
 Ser Ser Pro Gly Cys Gln Pro Pro Ala Gln Arg Leu Cys Gln Pro Val  
 530 535 540  
 Leu Pro Ser Pro Ala Cys Gln Leu Val Leu His Gln Ile Leu Lys Gly  
 545 550 555 560  
 Gly Ser Gly Thr Tyr Cys Leu Asn Val Ser Leu Ala Asp Thr Asn Ser  
 565 570 575  
 Leu Ala Val Val Ser Thr Gln Leu Ile Met Pro Gly Gln Glu Ala Gly  
 580 585 590  
 Leu Gly Gln Val Pro Leu Ile Val Gly Ile Leu Leu Val Leu Met Ala  
 595 600 605  
 Val Val Leu Ala Ser Leu Ile Tyr Arg Arg Arg Leu Met Lys Gln Asp  
 610 615 620  
 Phe Ser Val Pro Gln Leu Pro His Ser Ser Ser His Trp Leu Arg Leu  
 625 630 635 640  
 Pro Arg Ile Phe Cys Ser Cys Pro Ile Gly Glu Asn Ser Pro Leu Leu  
 645 650 655  
 Ser Gly Gln Gln Val Xaa  
 660

<210> 2  
 <211> 4  
 <212> PRT  
 <213> Mus musculus

<400> 2  
 Leu Glu Gly Ser  
 1

<210> 3  
 <211> 4  
 <212> PRT  
 <213> Mus musculus

<400> 3  
 Pro Lys Pro Pro  
 1

<210> 4  
 <211> 4  
 <212> PRT  
 <213> Mus musculus

<400> 4  
 Pro Leu Leu Pro  
 1

<210> 5  
 <211> 5  
 <212> PRT  
 <213> Mus musculus

<400> 5  
 Arg Ala Arg Gly Leu  
 1 5

<210> 6  
 <211> 23  
 <212> DNA  
 <213> Mus musculus

<400> 6  
 atgccccaaag aagacattca ctt

23

<210> 7  
 <211> 295  
 <212> DNA  
 <213> Mus musculus

<400> 7  
 tggttatccc aggaaggggc acagacgctc ctatgtcact gctgaagagg ccgcagggat 60  
 cggcatcctg atcgtggctc tggggattgc tctgcttata ggctgctggt actgtagaag 120  
 acgaagtgga tacagaacct tgatggacaa aaggcgctcat attggtattc aaaaaacctc 180  
 aagggaaaga tgcctcatgtg agagccctga tcaccaggac agccgactgt cttctcaaga 240  
 gaaatcccat cagcccgttg ttcccaacgc tccgcctgct tatgagaagc tctct 295

<210> 8  
 <211> 24  
 <212> DNA  
 <213> Mus musculus

<400> 8  
 tcaccgcccac cttattcacc ctga

24

<210> 9  
 <211> 357  
 <212> DNA  
 <213> Homo sapiens

<400> 9  
 atgccaagag aagatgctca cttcatctat ggttacccca agaaggggca cggccactct 60  
 tacaccacgg ctgaagaggg cgtctgggac ggcacccctga cagtgatccr gggagtctta 120  
 ctgctcatcg gctgttggga ttgtagaaga cgaaatggat acagagccct gatggataaa 180  
 agtcttcatg ttggcactca atgtgcctta acaagaagat gccacaaga agggtttgat 240  
 catcggggaca gcaaagtgtc tcttcaagag aaaaactgtg aacctgtggt tcccaatgct 300

ccaccctgccc atgagaaact ctctgcagaa cagucaccac caccctattc accttaa 357

<210> 10

<211> 8

<212> PRT

<213> Mus musculus

<400> 10

Met Pro Gln Glu Asp Ile His Phe  
1 5

<210> 11

<211> 98

<212> PRT

<213> Mus musculus

<400> 11

Gly Tyr Pro Arg Lys Gly His Arg Arg Ser Tyr Val Thr Ala Glu Glu  
1 5 10 15

Ala Ala Gly Ile Gly Ile Leu Ile Val Val Leu Gly Ile Ala Leu Leu  
20 25 30

Ile Gly Cys Trp Tyr Cys Arg Arg Arg Ser Gly Tyr Arg Thr Leu Met  
35 40 45

Asp Lys Arg Arg His Ile Gly Ile Gln Lys Thr Ser Arg Glu Arg Cys  
50 55 60

Ser Cys Glu Ser Pro Asp His Gln Asp Ser Arg Leu Ser Ser Gln Glu  
65 70 75 80

Lys Ser His Gln Pro Val Val Pro Asn Ala Pro Pro Ala Tyr Glu Lys  
85 90 95

Leu Ser

<210> 12

<211> 8

<212> PRT

<213> Mus musculus

<400> 12

Ser Pro Pro Pro Tyr Ser Pro Xaa  
1 5

<210> 13

<211> 119

<212> PRT

<213> Homo sapiens

<400> 13

Met Pro Arg Glu Asp Ala His Phe Ile Tyr Gly Tyr Pro Lys Lys Gly  
1 5 10 15

His Gly His Ser Tyr Thr Thr Ala Glu Glu Ala Ala Gly Ile Gly Ile

20 25 30  
 Leu Thr Val Ile Leu Gly Val Leu Leu Leu Ile Gly Cys Trp Tyr Cys  
 35 40 45  
 Arg Arg Arg Asn Gly Tyr Arg Ala Leu Met Asp Lys Ser Leu His Val  
 50 55 60  
 Gly Thr Gln Cys Ala Leu Thr Arg Arg Cys Pro Gln Glu Gly Phe Asp  
 65 70 75 80  
 His Arg Asp Ser Lys Val Ser Leu Gln Glu Lys Asn Cys Glu Pro Val  
 85 90 95  
 Val Pro Asn Ala Pro Pro Ala Tyr Glu Lys Leu Ser Ala Glu Gln Ser  
 100 105 110  
 Pro Pro Pro Tyr Ser Pro Xaa  
 115

<210> 14  
 <211> 1559  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (54)..(407)

<400> 14  
 agcagacaga ggaactctcat taaggaaggt gtctgtgtgcc ctgacctac aag atg 56  
 Met  
 1  
 cca aga gaa gat gct cac ttc atc tat ggt tac ccc aag aag ggg cac 104  
 Pro Arg Glu Asp Ala His Phe Ile Tyr Gly Tyr Pro Lys Lys Gly His  
 5 10 15  
 ggc cac tct tac acc acg gct gaa gag gcc gct ggg atc ggc atc ctg 152  
 Gly His Ser Tyr Thr Thr Ala Glu Glu Ala Ala Gly Ile Gly Ile Leu  
 20 25 30  
 aca gtg atc ctg gga gtc tta ctg ctc atc ggc tgt tgg tat tgt aqa 200  
 Thr Val Ile Leu Gly Val Leu Leu Leu Ile Gly Cys Trp Tyr Cys Arg  
 35 40 45  
 aga cga aat gga tac aga gcc ttg atg gat aaa agt ctt cat gtt ggc 248  
 Arg Arg Asn Gly Tyr Arg Ala Leu Met Asp Lys Ser Leu His Val Gly  
 50 55 60 65  
 act caa tgt gcc tta aca aga aga tgc cca caa gaa ggg ttt gat cat 296  
 Thr Gln Cys Ala Leu Thr Arg Arg Cys Pro Gln Glu Gly Phe Asp His  
 70 75 80  
 cgg gac agc aaa gtg tct ctt caa gag aaa aac tgt gaa cct gtg gtt 344  
 Arg Asp Ser Lys Val Ser Leu Gln Glu Lys Asn Cys Glu Pro Val Val  
 85 90 95  
 ccc aat gct cca cct gct tat gag aaa ctc tct gca gaa caa tca cca 392  
 Pro Asn Ala Pro Pro Ala Tyr Glu Lys Leu Ser Ala Glu Gln Ser Pro

100 105 110

cca cct tat tca cct taagagccag ccagacacct gagacatgct gaaattattt 447  
 Pro Pro Tyr Ser Pro  
 115

ctctcacact ttgcttgaa tttaatacag acatctaattg ttctcctttg gaatgggtga 507  
 ggaaaaatgc aagccatctc taataataag tcagtgttaa aatttttagta ggtccgctag 567  
 cagtactaat catgtgagga aatgatgaga aatattaaat tgggaaaact ccatacaataa 627  
 atgttgcaat gcatgatact atctgtgccg gaggtaatgt tagtaaatcc atggtgttat 687  
 tttctgagag acagaattca agtgggtatt cgggggcat ccaatttctc ttactttgaa 747  
 atttggctaa taacaaaacta gtcaggtttt cgaaccttga ccgacatgaa ctgtacacag 807  
 aattgttcca gtactatgga gtgctcacia aggatacttt tacagggttaa gacaaagggc 867  
 tgactggcct atttatctga tcaagaacat gtcagcaatg cctctttgtg ctctaaaatt 927  
 ctattatact acaataatat attgtaaaga tcttatagct cttttttttt gagatggagt 987  
 ttogcttttg ttgccaggc tggagtgcaa tggcgcgctc ttggctcacc ataacctccg 1047  
 cctcccaggt tcaagcaatt ctctgcctt agcctcctga gtagctggga ttacaggcgt 1107  
 gcgcactat gctgactaa tttgtagtt ttagtagaga cgggggttct ccatgttggt 1167  
 caggctggtc tcaaaactct gacctcaggt gatctgccc cctcagcctc ccaaagtgt 1227  
 ggaattacag gctgagcca ccacgcctgg ctggatccta tatcttaggt aagacatata 1287  
 aogcagtcta attacatttc actcaaggc tcaatgctat tctaactaat gacaagtatt 1347  
 ttctactaaa ccagaaattg gtagaaggat ttaataagt aaaagctant atgtaactgc 1407  
 ttagtgctga tgcctgtgta ctgccttaaa tgtacctatg gcaatttagc tctcttgggt 1467  
 tcccaaatcc ctctcacaag aatgtgcaga agaaatcata aaggatcaga gattctgaaa 1527  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 1559

&lt;210&gt; 15

&lt;211&gt; 118

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

Met Pro Arg Glu Asp Ala His Phe Ile Tyr Gly Tyr Pro Lys Lys Gly  
 1 5 10 15

His Gly His Ser Tyr Thr Thr Ala Glu Glu Ala Ala Gly Ile Gly Ile  
 20 25 30

Leu Thr Val Ile Leu Gly Val Leu Leu Leu Ile Gly Cys Trp Tyr Cys  
 35 40 45

Arg Arg Arg Asn Gly Tyr Arg Ala Leu Met Asp Lys Ser Leu His Val

50                      55                      60  
 Gly Thr Gln Cys Ala Leu Thr Arg Arg Cys Pro Gln Glu Gly Phe Asp  
 65                      70                      75                      80  
 His Arg Asp Ser Lys Val Ser Leu Gln Glu Lys Asn Cys Glu Pro Val  
 85                      90                      95  
 Val Pro Asn Ala Pro Pro Ala Tyr Glu Lys Leu Ser Ala Glu Gln Ser  
 100                      105                      110  
 Pro Pro Pro Tyr Ser Pro  
 115

<210> 16  
 <211> 517  
 <212> PRT  
 <213> Mus Musculus

<400> 16  
 Met Gly Leu Val Gly Trp Gly Leu Leu Leu Gly Cys Leu Gly Cys Gly  
 1                      5                      10                      15  
 Ile Leu Leu Arg Ala Arg Ala Gln Phe Pro Arg Val Cys Met Thr Leu  
 20                      25                      30  
 Asp Gly Val Leu Asn Lys Glu Cys Cys Pro Pro Leu Gly Pro Glu Ala  
 35                      40                      45  
 Thr Asn Ile Cys Gly Phe Leu Gln Gly Arg Gly Gln Cys Ala Glu Val  
 50                      55                      60  
 Gln Thr Asp Thr Arg Pro Trp Ser Gly Pro Tyr Ile Leu Arg Asn Gln  
 65                      70                      75                      80  
 Asp Asp Arg Glu Gln Trp Pro Arg Lys Phe Phe Asn Arg Thr Cys Lys  
 85                      90                      95  
 Cys Thr Gly Asn Phe Ala Gly Tyr Asn Cys Gly Gly Cys Lys Phe Gly  
 100                      105                      110  
 Trp Thr Gly Pro Asp Cys Asn Arg Lys Lys Pro Ala Ile Leu Arg Arg  
 115                      120                      125  
 Asn Ile His Ser Leu Thr Ala Gln Glu Arg Glu Gln Phe Leu Gly Ala  
 130                      135                      140  
 Leu Asp Leu Ala Lys Lys Ser Ile His Pro Asp Tyr Val Ile Thr Thr  
 145                      150                      155                      160  
 Gln His Trp Leu Gly Leu Leu Gly Pro Asn Gly Thr Gln Pro Gln Ile  
 165                      170                      175  
 Ala Asn Cys Ser Val Tyr Asp Phe Phe Val Trp Leu His Tyr Tyr Ser  
 180                      185                      190  
 Val Arg Asp Thr Leu Leu Gly Pro Gly Arg Pro Tyr Lys Ala Ile Asp  
 195                      200                      205  
 Phe Ser His Gln Gly Pro Ala Phe Val Thr Trp His Arg Tyr His Leu

210                      215                      220  
 Leu Trp Leu Glu Arg Glu Leu Gln Arg Leu Thr Gly Asn Glu Ser Phe  
 225                      230                      235                      240  
 Ala Leu Pro Tyr Trp Asn Phe Ala Thr Gly Lys Asn Glu Cys Asp Val  
                     245                      250                      255  
 Cys Thr Asp Asp Trp Leu Gly Ala Ala Arg Gln Asp Asp Pro Thr Leu  
                     260                      265                      270  
 Ile Ser Arg Asn Ser Arg Phe Ser Thr Trp Glu Ile Val Cys Asp Ser  
                     275                      280                      285  
 Leu Asp Asp Tyr Asn Arg Arg Val Thr Leu Cys Asn Gly Thr Tyr Glu  
                     290                      295                      300  
 Gly Leu Leu Arg Arg Asn Lys Val Gly Arg Asn Asn Glu Lys Leu Pro  
 305                      310                      315                      320  
 Thr Leu Lys Asn Val Gln Asp Cys Leu Ser Leu Gln Lys Phe Asp Ser  
                     325                      330                      335  
 Pro Pro Phe Phe Gln Asn Ser Thr Phe Ser Phe Arg Asn Ala Leu Glu  
                     340                      345                      350  
 Gly Phe Asp Lys Ala Asp Gly Thr Leu Asp Ser Gln Val Met Asn Leu  
                     355                      360                      365  
 His Asn Leu Ala His Ser Phe Leu Asn Gly Thr Asn Ala Leu Pro His  
                     370                      375                      380  
 Ser Ala Ala Asn Asp Pro Val Phe Val Val Leu His Ser Phe Thr Asp  
 385                      390                      395                      400  
 Ala Ile Phe Asp Glu Trp Leu Lys Arg Asn Asn Pro Ser Thr Asp Ala  
                     405                      410                      415  
 Trp Pro Gln Glu Leu Ala Pro Ile Gly His Asn Arg Met Tyr Asn Met  
                     420                      425                      430  
 Val Pro Phe Phe Pro Pro Val Thr Asn Glu Glu Leu Phe Leu Thr Ala  
                     435                      440                      445  
 Glu Gln Leu Gly Tyr Asn Tyr Ala Val Asp Leu Ser Glu Glu Glu Ala  
                     450                      455                      460  
 Pro Val Trp Ser Thr Thr Leu Ser Val Val Ile Gly Ile Leu Gly Ala  
 465                      470                      475                      480  
 Phe Val Leu Leu Leu Gly Leu Leu Ala Phe Leu Gln Tyr Arg Arg Leu  
                     485                      490                      495  
 Arg Lys Gly Tyr Ala Pro Leu Met Glu Thr Gly Leu Ser Ser Lys Arg  
                     500                      505                      510  
 Tyr Thr Glu Glu Ala  
                     515

&lt;210&gt; 17

&lt;211&gt; 519

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

```

Met Ser Pro Leu Trp Trp Gly Phe Leu Leu Ser Cys Leu Gly Cys Lys
 1          5          10          15

Ile Leu Pro Gly Ala Gln Gly Gln Phe Pro Arg Val Cys Met Thr Val
 20          25          30

Asp Ser Leu Val Asn Lys Glu Cys Cys Pro Arg Leu Gly Ala Glu Ser
 35          40          45

Ala Asn Val Cys Gly Ser Gln Gln Gly Arg Gly Gln Cys Thr Glu Val
 50          55          60

Arg Ala Asp Thr Arg Pro Trp Ser Gly Pro Tyr Ile Leu Arg Asn Gln
 65          70          75          80

Asp Asp Arg Glu Leu Trp Pro Arg Lys Phe Phe His Arg Thr Cys Lys
 85          90          95

Cys Thr Gly Asn Phe Ala Gly Tyr Asn Cys Gly Asp Cys Lys Phe Gly
100          105          110

Trp Thr Gly Pro Asn Cys Glu Arg Lys Lys Pro Pro Val Ile Arg Gln
115          120          125

Asn Ile His Ser Leu Ser Pro Gln Glu Arg Glu Gln Phe Leu Gly Ala
130          135          140

Leu Asp Leu Ala Lys Lys Arg Val His Pro Asp Tyr Val Ile Thr Thr
145          150          155          160

Gln His Trp Leu Gly Leu Leu Gly Pro Asn Gly Thr Gln Pro Gln Phe
165          170          175

Ala Asn Cys Ser Val Tyr Asp Phe Phe Val Trp Leu His Tyr Tyr Ser
180          185          190

Val Arg Asp Thr Leu Leu Gly Pro Gly Arg Pro Tyr Arg Ala Ile Asp
195          200          205

Phe Ser His Gln Gly Pro Ala Phe Val Thr Trp His Arg Tyr His Leu
210          215          220

Leu Cys Leu Glu Arg Asp Leu Gln Arg Leu Ile Gly Asn Glu Ser Phe
225          230          235          240

Ala Leu Pro Tyr Trp Asn Phe Ala Thr Gly Arg Asn Glu Cys Asp Val
245          250          255

Cys Thr Asp Gln Leu Phe Gly Ala Ala Arg Pro Asp Asp Pro Thr Leu
260          265          270

Ile Ser Arg Asn Ser Arg Phe Ser Ser Trp Glu Thr Val Cys Asp Ser
275          280          285

Leu Asp Asp Tyr Asn His Leu Val Thr Leu Cys Asn Gly Thr Tyr Glu
290          295          300

```



Gly Leu Leu Arg Arg Asn Gln Met Gly Arg Asn Ser Met Lys Leu Pro  
 305 310 315 320  
 Thr Leu Lys Asp Ile Arg Asp Cys Leu Ser Leu Gln Lys Phe Asp Asn  
 325 330 335  
 Pro Pro Phe Phe Gln Asn Ser Thr Phe Ser Phe Arg Asn Ala Leu Glu  
 340 345 350  
 Gly Phe Asp Lys Ala Asp Gly Thr Leu Asp Ser Gln Val Met Ser Leu  
 355 360 365  
 His Asn Leu Val His Ser Phe Leu Asn Gly Thr Asn Ala Leu Pro His  
 370 375 380  
 Ser Ala Ala Asn Asp Pro Ile Phe Val Val Leu His Ser Phe Thr Asp  
 385 390 395 400  
 Ala Ile Phe Asp Glu Trp Met Lys Arg Phe Asn Pro Pro Ala Asp Ala  
 405 410 415  
 Trp Pro Gln Glu Leu Ala Pro Ile Gly His Asn Arg Met Tyr Asn Met  
 420 425 430  
 Val Pro Phe Phe Pro Pro Val Thr Asn Glu Glu Leu Phe Leu Thr Ser  
 435 440 445  
 Asp Gln Leu Gly Tyr Ser Tyr Ala Ile Asp Leu Pro Val Ser Val Glu  
 450 455 460  
 Glu Thr Pro Gly Trp Pro Thr Thr Leu Leu Val Val Met Gly Thr Leu  
 465 470 475 480  
 Val Ala Leu Val Gly Leu Phe Val Leu Leu Ala Phe Leu Gln Tyr Arg  
 485 490 495  
 Arg Leu Arg Lys Gly Tyr Thr Pro Leu Met Glu Thr His Leu Ser Ser  
 500 505 510  
 Lys Arg Tyr Thr Glu Glu Ala  
 515

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/06034

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC(6) : Please See Extra Sheet.		
US CL : Please See Extra Sheet.		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
U.S. : 424/184.1, 277.1, 287.1, 93.2, 93.71; 435/372, 373; 514/2, 44; 530/350, 806, 827, 828; 536/23.1, 23.5		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
DIALOG medicine index, APS, WEST		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	LIU, M. Transfected human dendritic cells as cancer vaccines. Nat. Biotech. April 1988, Vol. 16, pages 335-336, see entire document.	1-23
A	DONNELLY, J.J. et al. DNA vaccines. Ann. Rev. Immunol. 1997, Vol. 15, pages 617-648, see entire document.	1-23
A, P	PARDOLL, D.M. Cancer vaccines. Nature Medicine. May 1998, Vol. 4, No.5(suppl.), pages 525-531, see entire document.	1-23
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search	Date of mailing of the international search report	
29 JUNE 1999	03 AUG 1999	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer <i>D. Lawrence</i> F PIERRE VANDERVEGT	
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196	

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/06034

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BOCZKOWSKI, D. et al. Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. J. Exp. Med. 01 August 1996, Vol. 184, No. 2, pages 465-472, see entire document, abstract in particular.	1-3, 7-10, 14-17, 20, 22
X	US 5,679,647 A (CARSON et al) 21 October 1997, see entire document.	1-3, 13, 20-23
X, P ---- Y, P	US 5,844,075 A (KAWAKAMI et al) 01 December 1998, see entire document.	1, 13, 20, 22-23 ---- 2-12, 14-19, 21

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/06034

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A01N 37/18, 43/04, 63/00; A61K 31/70, 35/12, 35/36, 39/00; C12N 5/02, 5/06; C07K 1/00; C07H 21/02, 21/04

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/184.1, 277.1, 287.1, 93.2, 93.71; 435/372, 373; 514/2, 44; 530/350, 806, 827, 828; 536/23.1, 23.5